

An alternative distraction osteogenesis method for atrophic posterior mandible: Case report

 Kagan Deniz¹,  Sina Uckan²,  Firdevs Senel³

¹Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Baskent University, Ankara, Turkey

²Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Istanbul Medipol University, Istanbul, Turkey

³Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Beykent University, Istanbul, Turkey

Copyright@Author(s) - Available online at www.annalsmedres.org

Content of this journal is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.



Abstract

Vertical lengthening by distraction osteogenesis has been widely used for reconstruction of dentoalveolar defects. However if bone height between the alveolar crest and the anatomical structures is insufficient, performing an appropriate osteotomy and placement of distractors in desired position is impossible. In the presented case sagittal lengthening of the posterior mandible with distraction osteogenesis was achieved. After horizontal augmentation of the crest by intraoral cortical bone graft, 2 implants were inserted to the distraction area.

Keywords: Transport distraction osteogenesis; mandible; alveolar bone

INTRODUCTION

Insufficient alveolar ridge width and/or height may complicate the successful placement of dental implants. Autogenous bone grafting, distraction osteogenesis, guided bone regeneration, ridge splitting and bone spreading are commonly used surgical procedures to resolve this problem (1,2).

Among all these techniques autogenous bone grafting and alveolar distraction osteogenesis (ADO) are commonly applied procedures, generally with the aim of simplifying the placement of dental implants (1,3-5). ADO is effective for vertical ridge lengthening, autogenous bone grafting is effective for horizontal ridge widening (1).

In ADO, minimum 4-5 mm of bone height should be present above the osteotomy line for device fixation and avoiding avascular necrosis of the transport segment (3). If the height of bone is less than this amount, performing ADO for vertical lengthening may be impossible.

In this case report sagittal lengthening of the posterior mandible with distraction osteogenesis is presented.

CASE REPORT

A 53 year old woman with severe vertically and horizontally atrophic, partially edentulous mandible referred to the clinic

for dental implant insertion. The patient had no familial or systemic disease. Clinical and radiographic examination revealed that inferior alveolar nerve and mental foramen was so close to alveolar crest at the right mandibular posterior area and there was a periapical abscess on the right lateral incisor tooth (Figure 1). Lateral incisor was extracted and distraction was delayed for bone healing. Under general anesthesia with orotracheal intubation 2 cc Ultracaine DS Forte (Articaine HCl, 1/100.000 epinephrine) supraperiosteal injection was performed. Following the vestibular sulcular incision, mucoperiosteum was released and unidirectional alveolar distractor (KLS Martin, 20mm, Zurich Ramus Distractor, Tut-tlingen/Germany) was inserted. The vertical osteotomy line was designed parallel to the axis of lateral incisor and the horizontal osteotomy line was designed parallel to the alveolar crest on the same level with the mental foramen (Figure 2). After 1 week of latency period, sagittal distraction of the transport alveolar segment was initiated. The device was activated 0, 50 x 2 times a day to achieve 1 mm of advancement. On the 20th day of the distraction period, alveolar segment was transported 20 mm posteriorly (Figure 2-3-4). After 3 months of consolidation, the distractor was removed (Figure 5) and a horizontal ridge augmentation with mandibular ramus graft was performed. Mild paresthesia due to mental nerve damage was observed. Six months

Received: 30.01.2020 **Accepted:** 27.07.2020 **Available online:** 22.02.2021

Corresponding Author: Kagan Deniz, Faculty of Dentistry, Baskent University, Ankara, Turkey

E-mail: drkagandeniz@gmail.com

later, implants were inserted to augmented bone (Figure 4) and it was noticed that sensation of the lower lip was normal.




Figure 1. Radiographic view of the atrophic mandible before distraction osteogenesis




Figure 2. Diagram of the osteotomy design and distraction




Figure 3. Radiographic view of the distractor and transport segment during consolidation period




Figure 4. Intraoral view of the distractor and transport segment during consolidation period



Figure 5. Intraoral view of the distraction area after removal of the distractor




Figure 6. Placement of dental implants

DISCUSSION

Alveolar bone is one of the most difficult areas of the human body to build, because both hard and soft tissues are missing in a wet and mobile area. Alveolar distraction osteogenesis, intraoral or extraoral autogenous bone grafting and guided bone regeneration are commonly performed as surgical procedures to reconstruct the insufficient alveolar ridge height. The volume of the bone maintained in guided bone regeneration is limited. Autogenous bone grafting is a widely accepted treatment modality for the surgical repair of alveolar defects. The technique requires a second surgical site and an adequate soft tissue, for the complete closure (1,6).

Intraoral bone grafting is usually preferred for small defects, and autogenous bone can be harvested easily from the ramus, symphysis or tuber maxilla, despite the fact that their size will be limited (7). When large amounts of bone are needed, crista iliaca is mostly the preferred donor site. Although autogenous bone grafting is still the gold standard, morbidity of the donor site and resorption of the graft are the known problems. Additionally, graft failure because of the insufficient mucosal coverage is possible (1,8). For the larger defects; reconstruction with free flaps are treatment of choice; however, this technique is difficult and costly, and an experienced microsurgery team is needed (9).

Over the years the distraction osteogenesis method has been under development for vertical augmentation of the jaws prior to implant placement. However, if the bone over the anatomic structure is limited, performing the osteotomy lines and placement of the device is difficult and sometimes impossible. Sagittal placement of the distractor may be another treatment option. In the presented case horizontal and vertical osteotomy lines were performed away from the mental foramen. A semicircular transport segment on a level with mental foramen provided better vertical augmentation as the transport segment raised onto mental foramen. New bone formation over the mental foramen and inferior alveolar nerve was observed and soft tissues were reshaped according to this new bone.

A gap may develop between the transport segment and basal segment due to the intervening of the soft tissue between the segments, and this may result in need for secondary grafting (10).

In our case although desired alveolar bone height was obtained, the alveolar ridge width was still insufficient for implant insertion. Onlay bone graft harvested from right mandibular ramus was inserted 3 months following the end of distraction.

Vector control is one of the main issues to be considered during ADO. Basa et al (3) had treated a mandibular defect using a Liou cleft distractor and transported the dentoalveolar segment in the sagittal plane from anterior to posterior. They used an acrylic splint to prevent lingual tilting of the transport segment.

In the presented case transport segment moved easily to desired location without any displacement and sufficient bone height was obtained without any complication.

Mandibular alveolar transport segment distraction is a convenient treatment modality for reconstruction of mandibular posterior alveolar defects. It is recommended for large alveolar defects, for defects that cannot be treated by grafting methods, and for patients who do not accept extraoral bone grafting. In this method hard and soft tissue regeneration is satisfactory. However the necessity for patient adaptation, device cost and insufficient final bone width are the disadvantages of this technique (11).

Acknowledgement

The authors thank Dr Kubilay Işık for the illustrations.

Conflict of interest : The authors declare that they have no competing interest.

Financial Disclosure: There are no financial supports.

Patient informed consent: Patient informed consent: Written informed consent was given by the patient's parents.

REFERENCES

1. Uckan S, Veziroglu F, Dayangac E. Alveolar distraction osteogenesis versus autogenous onlay bone grafting for alveolar ridge augmentation: Technique, complications, and implant survival rates. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008;106:511-5.
2. Chiapasco M, Zaniboni M, Boisco M. Augmentation procedures for the rehabilitation of deficient edentulous ridges with oral implants. *Clin Oral Implants Res* 2006;17 Suppl 2:136-59.
3. Basa S, Varol A, Yilmaz S. Transport distraction osteogenesis of a dentoalveolar segment in the posterior mandible: a technical note. *J Oral Maxillofac Surg* 2007;65:1862-4.
4. Enislidis G, Fock N, Millesi-Schobel G, et al. Analysis of complications following alveolar distraction osteogenesis and implant placement in the partially edentulous mandible. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2005;100:25-30.
5. Mohanty R, Kumar N, Ravindran C. Vertical alveolar ridge augmentation by distraction osteogenesis. *J Clin Diagn Res* 2015;9:43-6
6. Pektas ZO, Kircelli BH, Bayram B, et al. Alveolar cleft closure by distraction osteogenesis with skeletal anchorage during consolidation. *Int J Oral Maxillofac*

- Implants 2008;23:147-52.
7. Zhao K, Wang F, Huang W, et al. Comparison of dental implant performance following vertical alveolar bone augmentation with alveolar distraction osteogenesis or autogenous onlay bone grafts: a retrospective cohort study. *J Oral Maxillofac Surg* 2017;75:2099-114.
 8. Sbordone C, Toti P, Guidetti F, et al. Volume changes of iliac crest autogenous bone grafts after vertical and horizontal alveolar ridge augmentation of atrophic maxillas and mandibles: a 6- year computerized tomographic follow-up. *J Oral Maxillofac Surg* 2012;70:2559-65.
 9. Christiano JG, Gaggl A, Burger H, et al. The microvascular corticocancellous femur flap for reconstruction of the anterior maxilla in adult cleft lip, palate and alveolus patients. *Cleft Palate Craniofac J* 2012;49:305-13.
 10. Giotakis N, Narayan B, Nayagam S. Distraction osteogenesis and nonunion of the docking site: is there an ideal treatment option. *Injury* 2007;38:100-7
 11. Uckan S, Senol G, Ogut E, et al. Horizontal alveolar transport distraction osteogenesis followed by implant placement. *J Oral Maxillofac Surg* 2019;48:824-9.